

## Ginseng: a potentially effective Tcm for Alzheimer's disease

Na Li<sup>1</sup>, Yuan Liu<sup>1</sup>, Wenhao Yan<sup>3</sup>, Xin Sui<sup>1</sup>, Hui Li<sup>2</sup>, Xiaohua Li<sup>1</sup>, Xiaozheng Shi<sup>1</sup>, Qing Yang<sup>1</sup>, Jiaming Sun<sup>1</sup>, Xingcheng Lan<sup>1</sup>, Xijun Chen<sup>1</sup>, Xiaobo Qu<sup>1</sup>, Jianan Lin<sup>1</sup>, Dianjun Sui<sup>1\*</sup>

<sup>1</sup>Jilin Provincial Key Laboratory of BioMacromolecules of Chinese Medicine, Jilin Ginseng Academy, Changchun University of Chinese Medicine, Changchun, China.

<sup>2</sup>Qianwei Hospital of Jilin Province, Changchun 130012, China

<sup>3</sup>Frontage Laboratories Co., Ltd, Shanghai 201203, China

**Abstract:** Aging is a serious problem which is challenging the entire human race. With the longer life of people, more and more people are suffering from Alzheimer's disease (AD), which is suffering from not only for the patients themselves but also for the society. To be the king herb of Traditional Chinese Medicine (TCM), ginseng had been used as a treatment for dementia with a long history. Based on the traditional effect of "Reinforcing Intelligence" of ginseng, we reorganized the literatures and references about ginseng on the treatment of AD.

**Keywords:** Ginseng; Alzheimer's disease; Traditional Chinese medicine

Received 24 June 2018, Revised 21 August 2018, Accepted 25 August 2018

\*Corresponding Author: Dianjun Sui, lhaln@hotmail.com

### 1. Introduction

Alzheimer's disease (AD) is the most common dementia, which is a chronic neurodegenerative disease. The mechanism of AD is not well understood because of complex pathogenic factors. And with the progress of disease pathology, it usually gets deterioration. With the progress of AD, it's clinical symptoms will occur in inarticulate speech, disorientation, mood swings, loss of motivation, not managing self-care and abnormal behavior[1-2]. Now there is no certain well effective treatment to AD, but TCM could slow down the progression of disease, especially ginseng has antioxidant, protects the damage neuron cell and has been used in treating AD for thousands of years in clinical treatment.

What is about of the treatment on AD?

Base on the fundamental theory of traditional Chinese medicine, Qi and Blood are the fundamental substances constituting of the human body, which could maintain life activities. The original Yang of the body can not only promote hemokinesis, but also warm the body. If there are troubles about renal function in warming and moving (Qi Hua) such as fatigue, amnesia, cold-limbs, soreness of waist, dry eyes, urorrhagia and other clinical manifestations would lead into occurrence of disease. With the aging process, deficiency of Qi and Blood, disharmony between Ying and Wei, five zang-organs imbalances can all lead to the clinical manifestations of Qi Stagnation and Blood Stasis. Then the lucid Yang fails to rise and the turbid Yin fails to descend. When the body comes to this pathology phase, the mind has been damaged, which is Traditional Chinese Medicine explains the occurrence of Alzheimer's disease.

TCM clinical AD belongs to the category of

dementia and amnesia. The lesion location of Alzheimer's disease is brain and also is closely relate to the kidney, heart, spleen and liver. The key organ of body focus on the kidney. TCM about the basic pathological factors of AD are phlegm, dampness and stasis which has three factors influence each other. TCM usually use the treatment like removing phlegm and blood stasis, nourishing heart, tonifying spleen and restoring normal coordination between heart and kidney as adjuvant therapies. There are many kinds of traditional Chinese medicine, such as ginseng, hookwood, gardenia, ginkgo and so on, which can treat Alzheimer's disease[3-5].

What is Ginseng ?

As the king herb of TCM, Ginseng is the slow-growing perennial plants with fleshy roots, belonging to the genus *Panax* of the family Araliaceae. Ginseng mostly grows up in the northeast of China and North America, typically in cooler climates areas. The chemical components of ginseng is more rich and more high quality especially in Jilin province.

The root of ginseng is most available in dried form, either whole or sliced. Its leaf is rich in some saponins which could sometimes also used in saponins extraction and separation. Folk medicine attributes various benefits to oral use of American ginseng and Asian ginseng (*P. ginseng*) roots. It also has promoting aphrodisiac, strength the energy, enhancing immunity, helping type II diabetes treatment, or the treatment for sexual dysfunction in men[6].

Ginseng may be used in the health food or drug such as in energy drinks, herbal teas[7], ginseng coffee[8]. It has effects on some hairdressing and cosmetics supplies such as in hair tonics and cosmetic preparations, etc.

The pharamology of the Ginseng

People have used ginseng to keep healthy for

many years in Asian countries. Researchers have shown that ginseng has certain effects on central nervous system, hematopoiesis, metabolism, platelet, blood pressure and so on. More and more evidence has shown that Panax ginseng can improve the function of learning and memory[9-11]. Recently researches indicate that ginseng has shown well effects on the protecting the neuron cell and the treatment of AD, which's mechanism is related to anti-inflammation, antioxidant, anti-apoptosis and anti-aging, etc.

## 2. Anti-amyloid A (A $\beta$ ) aggregation

A $\beta$  is the main pathogenesis-related proteins of AD according to recently researches. As reported, A $\beta$  concentration in neuron cell is obvious reduction after taking oral ginseng[12]. Zhang et al. reported that ginseng could alleviate A $\beta$  aggregation in ovariectomized (OVX) and D-galactose-injected rats by decreasing  $\beta$ -secretase and upregulating  $\alpha$ -secretase[13]. Ginseng Rg1 can inhibit the aggregation of A $\beta$  by up-regulate the expression of PPAR $\gamma$  and increase the content of IDE in hippocampus. Hippocampal histopathological abnormalities can improve learning and memory in a rat model of AD[14]. Ginsenoside Rd attenuates A $\beta$ -induced pathological phosphorylation by affecting the activities of GSK-3 $\beta$  and PP-2A[15]. Ginseng Rh2 treatment increases soluble APP $\alpha$  (sAPP $\alpha$ ) levels and reduces amyloid beta 40 and 42 concentrations by increasing CTF $\alpha$ / $\beta$  ratios[16]. In vitro, it had proved that ginseng could inhibit  $\beta$ -secretase activity and increase  $\alpha$ -secretase activity[17-18]. It also reported that ginseng could inhibit the activity of  $\gamma$ -secretase in TgmApp mice and B103-APP cells[19].

## 3. Anti-apoptosis

The apoptosis of neuron cell is also a damage factor of AD[20]. Li et al. reported that ginseng could increase neuron viability and decrease lactate dehydrogenase[21]. Gong et al. used real-time RT-PCR, flow cytometry and western-blot to confirm the effect of ginseng on hippocampal neurons. The result showed that ginseng could increase neuronal viability, decrease LDH and reverse the apoptosis[22]. Another research showed that SHSY5Y cell apoptosis induced by MPP $^{+}$  could be inhibited by ginseng and the anti-apoptosis function may work through alleviating activities of intracellular reactive oxygen species(ROS) and C-jun N-terminal protein kinase(JNK) and subsequently intimating activation of caspase3[23].

## 4. Anti-oxidation and Anti-inflammation

Excessive oxidants and neuro-inflammation are two pathological alterations in AD. Microglia which is found to be a mediator in the neuronal immune

system, plays an important role of AD in many pathological processes[24-26]. It has been proved that many proinflammatory cytokines like IL-1 $\beta$ , IL-6 and TNF- $\alpha$  are released from activated microglia[27]. Recently research showed that the antioxidant effect of ginseng could protect the nervous system in vitro[28]. In vivo, it had been proved that ginseng could regulate immunity[29]. Ginseng Rg5 can decrease A $\beta$  deposits, improve STZ-induced learning and memory impairments in rats by increasing expressions of BDNF and IGF-1[30]. Compound K can overcome scopolamine-induced memory impairment by the induction of Nrf2-mediated antioxidant enzymes[31].

Clinical studies have shown that taking ginseng in young and middle-aged people can improve memory and cognitive function[32,33]. More and more researchers found that ginseng has certain effects on neurotransmitters, calcium channel inhibition, mitochondrial dysfunction, signaling pathways, etc.

Other Effects about Ginseng:

### 1. Effects on cardiovascular system

Ginseng has been proved hypotensive effect in hypertensive rats, and also in patients with essential hypertension. Rg1 and Rg3 can relax vascular smooth muscle and inhibit endothelin production and platelet aggregation.

### 2. Effects on central nervous system

The active ingredients in ginseng, such as Rb1, Rg1, can enhance the activity of the central nervous system. For instance, ginsenoside Rg1 can improve the whole process of memory. And ginsenoside Rb1 can only improve the acquisition and retention ability of memory but has nothing to do with the consolidated function of memory.

### 3. Effects on anti-tumor

Rh2 and Rg3 have inhibitory effects on breast, prostate, liver and small bowel tumors in animals and human trials. Ginseng can improve the protective effect on the human anti X-ray radiation. Long term use of Ginseng can reduce the incidence of lung, stomach, liver and colon cancer. The main mechanism of all above is to enhance immunity, inhibit tumor angiogenesis and induce tumor cell apoptosis.

4. Effects on immune system, inflammation and allergy

Rb1, Rg1, Rg3 can suppress cytokine production, the expression of COX-2 gene and the release of histamine. Polysaccharide can increase the production of interferon, enhance the activity of phagocytic cell, natural killer cells, B and T cells. Rg1 can increase the function of t-helper cells and improve the immune function of the elderly.

## 5. Conclusion

Ginseng has been proved to have certain effects in treating Alzheimer disease. There are hundreds of prescriptions in TCM about ginseng treating AD. But

some mechanism and treatment of AD is still a problem. We would try our best to resolve the problem of ginseng mechanism and finding out some certain kinds of chemical substances. Make ginseng a new therapeutic strategy in preventing and treatment for AD.

### Acknowledgements

This work was supported by the Jilin Provincial Science and Technology Department (20160101229JC) and the National Natural Science Foundation of China (81704001). We thank Prof. Zhao Da-Qing and Prof. Zhao Yu (Changchun University of Chinese Medicine) gave more technology supports.

### References

- [1] Burns A, Iliffe S. Alzheimer's disease[J]. *BMJ (Clinical research)* 2009, 338:158.
- [2] Dubois B, Feldman H H, Jacova C, et al. Research criteria for the diagnosis of Alzheimer's disease: revising the NINCDS-ADRDA criteria[J]. *Lancet Neurology*, 2007, 6(8):734-746.
- [3] KA Wesnes T, Ward A, McGinty O. The memory enhancing effects of a Ginkgo biloba/Panax ginseng combination in healthy middle-aged volunteers[J]. Springer-Verlag, 2000, 152:353-361.
- [4] Yang Y, Ji WG, Zhu ZR, et al. Rhynchophylline suppresses soluble A $\beta$ 1-42-induced impairment of spatial cognition function via inhibiting excessive activation of extrasynaptic NR2B-containing NMDA receptors[J]. *Neuropharmacology*, 2018, 3908(18):30108-30114.
- [5] Zang CX, Bao XQ, Li L, et al. The Protective Effects of Gardenia jasminoides (Fructus Gardenia) on Amyloid- $\beta$ -Induced Mouse Cognitive Impairment and Neurotoxicity[J]. *Am J Chin Med*, 2018, 46(2):1-17.
- [6] Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics[J]. *Science*, 2002, 297(5580):353-356.
- [7] Selkoe DJ. Alzheimer's Disease Is a Synaptic Failure[J]. *Science*, 2002, 298(5594):789-791.
- [8] Clauson KA, Shields KM, Mcqueen CE, et al. Safety issues associated with commercially available energy drinks[J]. *Journal of the American Pharmacists Association*, 2008, 48(3):55-67.
- [9] Cheng Y, Shen LH, Zhang JT. Anti-amnestic and anti-aging effects of ginsenoside Rg1 and Rb1 and its mechanism of action[J]. *Acta Pharmacologica Sinica*, 2005, 26(2):143-149.
- [10] Joo SS, Won TJ, Lee DI. Reciprocal activity of ginsenosides in the production of proinflammatory repertoire, and their potential roles in neuroprotection in vivo[J]. *Planta Medica*, 2005, 71(5):476-481.
- [11] Fang F, Chen X, Huang T, et al. Multi-faced neuroprotective effects of Ginsenoside Rg1 in an Alzheimer mouse model[J]. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 2012, 1822(2):286-292.
- [12] Chen F, Eckman E A, Eckman C B. Reductions in levels of the Alzheimer's amyloid beta peptide after oral administration of ginsenosides[J]. *Faseb Journal Official Publication of the Federation of American Societies for Experimental Biology*, 2006, 20(8):1269-71.
- [13] Zhang X, Wang J, Xing Y, et al. Effects of ginsenoside Rg1 or 17 $\beta$ estradiol on a cognitively impaired, ovariectomized rat model of Alzheimer's disease[J]. *Neuroscience*, 2012, 220:191-200.
- [14] Quan QK, Wang J, Li X, et al. Ginsenoside Rg1 Decreases A $\beta$ 1-42 Level by Upregulating PPAR $\gamma$  and IDE Expression in the Hippocampus of a Rat Model of Alzheimer's Disease[J]. *Plos One*, 2013, 8(3):e59155.
- [15] Ling L, Liu Z, Liu J, et al. Ginsenoside Rd attenuates beta-amyloid-induced tau phosphorylation by altering the functional balance of glycogen synthase kinase 3beta and protein phosphatase 2A[J]. *Neurobiology of Disease*, 2013, 54(6):320-328.
- [16] Efthimiopoulos S, Punj S, Manolopoulos V, et al. Intracellular cyclic AMP inhibits constitutive and phorbol ester-stimulated secretory cleavage of amyloid precursor protein.[J]. *Journal of Neurochemistry*, 1996, 67(2):872-875.
- [17] Wang YH, Du GH. Ginsenoside Rg1 inhibits beta-secretase activity in vitro and protects against Abeta-induced cytotoxicity in PC12 cells.[J]. *Journal of Asian Natural Products Research*, 2009, 11(7):604.
- [18] Shi C, Zheng DD, Fang L, et al. Ginsenoside Rg1 promotes nonamyloidogenic cleavage of APP via estrogen receptor signaling to MAPK/ERK and PI3K/Akt[J]. *Biochimica et Biophysica Acta (BBA) - General Subjects*, 2012, 1820(4):453-460.
- [19] Fang F, Chen X, Huang T, et al. Multi-faced neuroprotective effects of Ginsenoside Rg1 in an Alzheimer mouse model[J]. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 2012, 1822(2):286-292.
- [20] Li WZ, Li WP, Zhang W, et al. Protective Effect of Extract of Astragalus on Learning and Memory Impairments and Neurons Apoptosis Induced by Glucocorticoids in 12 - Month Male Mice[J]. *Anatomical Record*, 2011, 294(6):1003.
- [21] Li J, Zhang J. Inhibition of apoptosis by ginsenoside Rg1 in cultured cortical

- neurons[J]. *Chin Med J*, 1997, 110(7):535-539.
- [22] Gong L, Li SL, Li H, et al. Ginsenoside Rg1 protects primary cultured rat hippocampal neurons from cell apoptosis induced by  $\beta$ -amyloid protein[J]. *Pharmaceutical Biology*, 2011, 49(5):501-507.
- [23] Fang F, Chen XC, Zhu YG, et al. Ginsenoside Rg1 may protect SHSY5Y cells from apoptosis induced by MPP<sup>+</sup> through JNK way[J]. *Acta Pharmaceutica Sinica*, 2003, 38(3):176.
- [24] Cameron B, Landreth GE. Inflammation, Microglia and Alzheimer's Disease[J]. *Neurobiology of Disease*, 2010, 37(3):503.
- [25] Urte N, Neher JJ, Brown GC. Neuronal Death Induced by Nanomolar Amyloid  $\beta$  Is Mediated by Primary Phagocytosis of Neurons by Microglia[J]. *Journal of Biological Chemistry*, 2011, 286(46):39904-39913.
- [26] Mandrekar-Colucci S, Landreth GE. Microglia and inflammation in Alzheimer's disease[J]. *Cns & Neurological Disorders Drug Targets*, 2010, 9(2):156.
- [27] Orellana JA, Sáez PJ, Shoji KF, et al. Modulation of brain hemichannels and gap junction channels by pro-inflammatory agents and their possible role in neurodegeneration[J]. *Antioxidants & Redox Signaling*, 2009, 11(2):369-99.
- [28] Liu Q, Kou JP, Yu BY. Ginsenoside Rg1 protects against hydrogen peroxide-induced cell death in PC12 cells via inhibiting NF- $\kappa$ B activation[J]. *Neurochemistry International*, 2011, 58(1):119-125.
- [29] Joo SS, Won TJ, Lee DI. Reciprocal activity of ginsenosides in the production of proinflammatory repertoire, and their potential roles in neuroprotection in vivo[J]. *Planta Medica*, 2005, 71(5):476-81.
- [30] Chu S, Gu J, Feng L, et al. Ginsenoside Rg5 improves cognitive dysfunction and beta-amyloid deposition in STZ-induced memory impaired rats via attenuating neuroinflammatory responses[J]. *International Immunopharmacology*, 2014, 19(2):317-326.
- [31] Ji Y S, Ju SH, Oh J, et al. Neuroprotective and Cognition-Enhancing Effects of Compound K Isolated from Red Ginseng[J]. *Journal of Agricultural & Food Chemistry*, 2016, 64(14):2855.
- [32] Scholey A, Ossoukhova A, Owen L, et al. Effects of American ginseng (*Panax quinquefolius*) on neurocognitive function: an acute, randomised, double-blind, placebo-controlled, crossover study[J]. *Psychopharmacology*, 2010, 212(3):345-356.
- [33] Ossoukhova A, Owen L, Savage K, et al. Improved working memory performance following administration of a single dose of American ginseng (*Panax quinquefolius* L.) to healthy middle-age adults[J]. *Human Psychopharmacology Clinical & Experimental*, 2015, 30(2):108-122.